

**Acknowledgment.** We thank the donors of the Petroleum Research Fund, administered by the American Chemical Society, for a grant in support of this work.

**Registry No.** 1, 105282-83-5; 2, 105282-78-8; 3, 105282-79-9; 4, 105282-80-2; 5, 105282-81-3; 6, 105282-82-4; 1,1-dimethyl-2-tetralone, 20027-76-3; 3-bromo-1,1-dimethyl-2-naphthalenone, 105282-76-6; 3-bromo-2-chloro-1,1-dimethyl-1,2-dihydronaphthalene, 105282-77-7; 1,3-diphenylisobenzofuran, 5471-63-6; 4,4-dimethyl-2-tetralone, 83810-57-5.


## Methylidene Migratory Insertion into an Ru-H Bond

Emily A. Carter and William A. Goddard III\*

Contribution No. 7409  
Arthur Amos Noyes, Laboratory of Chemical Physics  
California Institute of Technology  
Pasadena, California 91125  
Received May 5, 1986


The migratory insertions of  $\text{CH}_x$  fragments into transition-metal-hydrogen and transition-metal-alkyl bonds have long been proposed as chain initiation and propagation steps in the Fischer-Tropsch synthesis of hydrocarbons.<sup>1</sup> Particularly for ruthenium, an effective heterogeneous catalyst for the production of high molecular weight polymethylenes,<sup>2</sup> there is strong indirect evidence that the chain growth mechanism involves methylidene insertion into growing alkyl chains.<sup>1,3</sup> Several experiments on homogeneous systems point to the facility of direct  $\text{CH}_2$  insertions into both M-H and M-R bonds.<sup>4</sup> Thorn and Tulip<sup>5a</sup> proposed that acidification of a hydrido hydroxymethyliridium complex proceeds via a hydridomethyleneiridium intermediate which undergoes  $\text{CH}_2$  insertion into the Ir-H bond to yield an iridium methyl complex. Upon hydrogen abstraction from mononuclear metal dimethyl complexes, Thorn and Tulip,<sup>5b</sup> as well as Cooper,<sup>6</sup> Maitlis,<sup>7</sup> and Werner,<sup>8</sup> have postulated the intermediacy of methyl methylidene metal complexes which insert  $\text{CH}_2$  into M-CH<sub>3</sub> and then  $\beta$ -hydride eliminate en route to the formation of ethylene hydride complexes. Thus these studies suggest that both the chain initiation and propagation steps in Fischer-Tropsch synthesis may be facile even at a single metal center.

As a model for these important elementary reactions, we have used ab initio quantum mechanical techniques to investigate the migratory insertion of  $\text{CH}_2$  into an adjacent Ru-H bond. To our knowledge, these calculations provide the first quantitative description of the energetics of such a reaction, including evaluations of both the activation barrier to insertion as well as the relative stabilities of the reactant and product. The reaction pathway is depicted below



where **1** is a model for 18-electron complexes such as  $(\text{C}_5\text{H}_5)_-$


- (1) Biloen, P.; Sachtler, W. M. H. *Adv. Catal.* **1981**, *30*, 165.  
(2) (a) Anderson, R. B. In *Catalysis*; Emmett, P. H., Ed.; Reinhold: New York, 1956; Vol. IV, pp 237-242. (b) Pichler, H.; Buffleb, H. *Brennst.-Chem.* **1940**, *21*, 257, 273, 285.  
(3) (a) Brady, R. C., III; Pettit, R. *J. Am. Chem. Soc.* **1980**, *102*, 6181. (b) *Ibid.* **1981**, *103*, 1287. (c) Baker, J. A.; Bell, A. T. *J. Catal.* **1982**, *78*, 165-181.  
(4) The first observation of general alkylidene insertions into M-R bonds was by: Sharp, P. R.; Schrock, R. R. *J. Organomet. Chem.* **1979**, *171*, 43.  
(5) (a) Thorn, D. L.; Tulip, T. H. *Organometallics* **1982**, *1*, 1580. (b) Thorn, D. L.; Tulip, T. H. *J. Am. Chem. Soc.* **1981**, *103*, 5984.  
(6) Hayes, J. C.; Pearson, G. D. N.; Cooper, N. J. *J. Am. Chem. Soc.* **1981**, *103*, 4648.  
(7) Isobe, K.; Andrews, D. G.; Mann, B. E.; Maitlis, P. M. *J. Chem. Soc., Chem. Commun.* **1981**, 809.  
(8) Kletzin, H.; Werner, H.; Serhadli, O.; Ziegler, M. L. *Angew. Chem., Int. Ed. Engl.* **1983**, *22*, 46.




**Figure 1.** Reaction coordinate for the insertion of  $\text{CH}_2$  into Ru-H in **1** to form  $\text{ClRu}(\text{CH}_3)$  (**2**) at the HF, GVB-PP(3/6), GVB-RCI(3/6), GVB-RCI(3/6), and GVB-RCI(3/6)-MCSCF levels. Energy (kcal/mol) is plotted relative to the total energy for **2** vs.  $R(\text{Ru-H})/[R(\text{Ru-H}) + R(\text{C-H})]$  (normalized reaction coordinate). Also shown at the top are the corresponding H-Ru-C angles (deg). The full GVB-RCI-MCSCF leads simultaneously to a proper description of both the reactant-like and product-like configurations important at the transition state and hence to a smooth potential curve. Some lower level calculations lead to a less smooth transition, the wave function being less capable of simultaneous description of both reactant and product channels.

$(\text{PPh}_3)_2\text{Ru}(\text{R})(\text{CH}_2)$  (**3**) or  $[(\text{C}_6\text{Me}_6)(\text{PPh}_3)_2\text{Ru}(\text{CH}_3)(\text{CH}_2)]^+$  (**4**), the intermediate postulated by Werner.<sup>8</sup> As discussed previously,<sup>9</sup> **1** conforms to the valence bond (VB) view of oxidation states in which electronegative ligands may remove no more than two units of charge from the metal (the easily ionized s-electrons), leaving the other ligands to form covalent bonds to unpaired d-electrons (or to form donor bonds into empty metal valence orbitals). Thus ligands with large electron affinities<sup>10</sup> such as Cp ( $\eta^5\text{-C}_5\text{H}_5$ ) and Cl form rather ionic bonds with the metal valence electrons, while neutral  $\pi$ -donor ligands (e.g.,  $\pi$ -aryls) and phosphines require empty metal valence orbitals. Finally, ligands with unpaired electrons (and small electron affinities, e.g.,  $\text{CH}_2$ ,  $\text{CH}_3$ , H, NO, etc.) require unpaired metal d-electrons with which to form covalent bonds. As a result, we believe the singlet state of **1** is a good model for **3** and **4**, since all three complexes have a metal VB oxidation state of +1. Ru(I) is  $d^7$ , with three unpaired d-electrons to form covalent bonds to R and  $\text{CH}_2$  in **1**, **3**, and **4**.

Consider the process of inserting the  $\text{CH}_2$  ligand into the Ru-H bond to form an Ru- $\text{CH}_3$  species. We begin with an Ru-H bond and two Ru-C in-plane bonds (one  $\sigma$  and one  $\pi$ ) which are converted to a C-H bond, one Ru-C bond and an Ru d lone pair. Notice that the presence of the in-plane  $\pi$ -bond<sup>11</sup> suggests a



- (9) Carter, E. A.; Goddard, W. A., III *J. Am. Chem. Soc.* **1986**, *108*, 2180.  
(10) The electron affinities of Cp and Cl are 2.2 eV (Rosenstock, H. M.; Draxl, K.; Steiner, B. W.; Herron, J. T. *J. Phys. Chem. Ref. Data* **1977**, *6*, 736-772) and 3.62 eV (Hotop, H.; Lineberger, W. C. *J. Chem. Phys. Ref. Data* **1975**, *4*, 539-576), respectively.  
(11) This conformation is the lowest energy orientation for **1**: Carter, E. A.; Goddard, W. A., III, manuscript in preparation.



**Figure 2.** GVB(3/6)PP one-electron orbitals near the transition state ( $\Theta(\text{H-Ru-C}) = 50.0^\circ$ ). (a) Orbital pair describing the Ru-H bond of the reactant **1** and the C-H bond of the product **2**; (b) orbital pair describing the Ru-C  $\sigma$ -bond for both **1** and **2**; (c) orbital pair describing the Ru-C  $\pi$ -bond of **1** and the Ru  $d\pi$  lone pair of **2**. Contours are shown at intervals of 0.05 au.

smooth transition from an Ru-H to a C-H bond may be possible, since the in-plane carbon p-orbital is oriented correctly for formation of the in-plane C-H bond. Indeed, at the highest level of theory examined, we find that the  $\text{CH}_2$  insertion into Ru-H proceeds with a low activation barrier (11.5 kcal/mol) and is thermodynamically favorable, with an exothermicity of 7.1 kcal/mol, as displayed in the reaction coordinate of Figure 1. Notice that the transition state occurs approximately halfway between reactants and products, as expected for a reaction which is nearly thermoneutral (Hammond postulate).

Figure 2 shows the orbitals near the transition state [ $\Theta(\text{H-Ru-C}) = 50^\circ$ ]. Here we see that the Ru-H bond smoothly converts into the C-H bond (Figure 2a), while the Ru-C  $\sigma$ -bond (Figure 2b) does not change significantly. At the transition state, the Ru-C  $\pi$ -bond (Figure 2c) has begun to move out of the way of the incipient C-H bond and already has substantial Ru  $d$  lone-pair character.<sup>12</sup> The Ru-C and Ru-H bonds at the transition state have lengthened significantly from their values in **1**, increasing from 1.87 to 1.93 Å for Ru-C and from 1.65 to 1.77 Å for Ru-H.

The exothermicity, activation barrier, and transition-state geometry were calculated at five levels of theory, as shown in Figure 1.<sup>13</sup> The geometries along the reaction coordinate were predicted by analytic gradients of Hartree-Fock wave functions,<sup>14</sup> with all geometrical parameters optimized at each H-Ru-C angle.<sup>15</sup> In

(12) By the point at which  $\Theta(\text{H-Ru-C}) = 40.0^\circ$ , the Ru  $d$  lone pair is fully formed. (The equilibrium geometry of  $\text{ClRuCH}_3$  has an H-Ru-C angle of  $23.2^\circ$ .)

(13) Full details to be published elsewhere. A valence double- $\zeta$  quality basis was used.

(14) Hartree-Fock (HF) calculations are known to predict accurate geometries. As a test, we optimized the geometry of **1** at both the HF and GVB-RCI(3/6) levels and found that the two geometries were very similar (e.g., all bond lengths and angles differed by at most 0.03 Å and  $11.9^\circ$ , respectively); see ref 11.

(15) The geometries of **1** and **2** were optimized with no restrictions except the retention of  $C_s$  symmetry (lower symmetry cases were found to be higher in energy; ref 11).

the highest level of theory considered (the bottom curve of Figure 1), we optimize the six active orbitals (the orbitals actively involved in the insertion, namely, the Ru-H and the Ru-C  $\sigma$ - and  $\pi$ -bond pairs) self-consistently for a full six-electron CI within those six orbitals (all occupations of six electrons in six orbitals—the GVB(3/6)CI-MCSCF level). This level allows a balanced description of the three bond pairs changing during the reaction. Higher level, extended basis dissociation-consistent CI calculations<sup>16</sup> on various dissociation processes involving these species<sup>13</sup> suggest that the true exothermicity is 10.4 kcal/mol, in good agreement with our MCSCF calculations.

In conclusion, we have shown that alkylidene migratory insertions can be quite facile, proceeding with a low activation barrier. These calculations provide the first quantitative evidence for the feasibility of this elementary reaction (previously postulated based on experimental results,<sup>4-8</sup> but never directly observed). These results suggest that for Ru, the reverse reaction of  $\alpha$ -hydrogen elimination is subject to a barrier of 18.6 kcal/mol. This is consistent with the fact that  $\alpha$ -H eliminations most often occur for the early transition metals. Work in progress on the related reaction of  $\text{CH}_2$  insertion into an Ru-alkyl bond suggests an exothermicity of 4.9 kcal/mol. The activation barrier will probably be higher than that for H due to the necessary reorientation of the alkyl upon migration from Ru to  $\text{CH}_2$ .<sup>17</sup> The alkyl migration differs primarily from the hydride energetics because the incipient C-C bond is weaker than the incipient C-H bond. While our calculations suggest that late transition metals undergo  $\text{CH}_2$  insertion with relative ease, early metal alkylidenes have been observed that do *not* insert into M-R bonds. We believe that this is due to the much greater strength of the M-C  $\pi$ -bond for the early transition metals.<sup>18</sup>

**Acknowledgment.** This work was supported by the Shell Development Co., Houston, TX, and the National Science Foundation (Grant CHE83-18041). E.A.C. gratefully acknowledges the support of a National Science Foundation Predoctoral Fellowship (1982-1985).

(16) Bair, R. A.; Goddard, W. A., III, unpublished results. Bair, R. A. Ph.D. thesis, Caltech, 1980. Carter, E. A.; Goddard, W. A., III *J. Phys. Chem.* **1984**, *88*, 1485. Reference 9.

(17) Low, J. J.; Goddard, W. A., III *Organometallics* **1986**, *5*, 609.

(18) Carter, E. A.; Goddard, W. A., III *J. Am. Chem. Soc.* **1986**, *108*, 4746.

## Carbyne-Carbyne Ligand Coupling on a Single Tungsten Center

Gregory A. McDermott and Andreas Mayr\*

Department of Chemistry, Princeton University  
Princeton, New Jersey 08544

Received September 10, 1986

The coupling of two carbyne ligands on a single transition-metal center has been analyzed theoretically by Hoffmann, Wilker, and Eisenstein.<sup>1</sup> These authors concluded that the reaction should be allowed for systems containing six electrons in the metal-ligand  $\pi$ -orbitals. In Fischer-type carbyne complexes, e.g.,  $[(\text{W} \equiv \text{CPh})\text{Br}(\text{CO})_4]$  (**1**), there are six electrons in metal-ligand  $\pi$ -orbitals.<sup>2</sup> The carbyne complex **1** is derived from tungsten hexacarbonyl, itself a six- $\pi$ -electron system, by transformation of a carbonyl ligand into a carbyne ligand. Since the number of  $\pi$ -electrons is not changed during this reaction, a second transformation of one of the remaining carbonyl ligands in **1** into a carbyne ligand should lead to a tungsten bis(carbyne) system with

(1) (a) Hoffmann, R.; Wilker, C. N.; Eisenstein, O. *J. Am. Chem. Soc.* **1982**, *104*, 632-634. (b) Wilker, C. N.; Hoffmann, R.; Eisenstein, O. *Nouv. J. Chim.* **1983**, *7*, 535-544.

(2) (a) Fischer, E. O.; Schubert, U. *J. Organomet. Chem.* **1975**, *100*, 59-81. (b) Kostič, N. M.; Fenske, R. F. *Organometallics* **1982**, *1*, 489-496.